




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : A61K 41/00		A2	(11) International Publication Number: WO 00/62808
			(43) International Publication Date: 26 October 2000 (26.10.00)
(21) International Application Number: PCT/US00/10100		(71)(72) Applicants and Inventors: COLLINS, Douglas, A. [US/US]; 1150 Meadowlark Court S.W., Rochester, MN 55902 (US). HOGENKAMP, Henricus, P., C. [US/US]; 2211 Marion Road, Roseville, MN 55113 (US).	
(22) International Filing Date: 15 April 2000 (15.04.00)		(74) Agent: VIKSNINS, Ann, S.; Schwegman, Lundberg, Woessner & Kluth, P.O. Box 2938, Minneapolis, MN 55402 (US).	
(30) Priority Data: 60/129,733 16 April 1999 (16.04.99) US 60/159,873 15 October 1999 (15.10.99) US		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications US 60/129,733 (CIP) Filed on 16 April 1999 (16.04.99) US 60/159,873 (CIP) Filed on 15 October 1999 (15.10.99)		Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(71) Applicants (for all designated States except US): MAYO OF FOUNDATION FOR MEDICAL EDUCATION AND RESEARCH [US/US]; 200 First Street S.W., Rochester, MN 55905 (US). REGENTS OF THE UNIVERSITY OF MINNESOTA [US/US]; 600 University Gateway, 200 Oak Street SE, Minneapolis, MN 55455-2070 (US).			

(54) Title: **COBALAMIN CONJUGATES USEFUL AS ANTITUMOR AGENTS**

(57) Abstract

The invention provides cobalamin compounds linked to a neutron capture therapy target (e.g. Boron-10 or Gadolinium-157), and optionally linked to a detectable moiety, as well as pharmaceutical compositions comprising the compounds, and methods for using the compounds in medical diagnosis and therapy.



BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Nishimura Co., Ltd (1986)) for the treatment of malignant gliomas and those of Mishima et al. (Mishima, Y. et al., *Lancet*, 2, 388-289 (1989)) for melanoma, there has been renewed national and international interest in BNCT.

5 The theoretical advantage of BNCT is that it is a two component or binary system, consisting of ^{10}B and thermal neutrons, which when combined together generate high linear energy transfer (LET) radiation capable of selectively destroying tumor cells without significant damage to normal tissues. In order for BNCT to succeed a critical amount of ^{10}B and a sufficient number of thermal neutrons must be delivered to individual tumor cells.

10 Over the past few years the Department of Energy and the NIH have renewed funding for BNCT-related research, and this has supported a growing number of investigators in many different disciplines. Advances in BNCT in the areas of compound distribution and pharmacokinetics compare favorably with other emerging modalities such as photon activation therapy, photodynamic therapy, and the use of radiolabeled antibodies for cancer
15 treatment in which physiological targeting is used.

There are a number of nuclides that have a high propensity for absorbing low energy or thermal neutrons, and this property, referred to as the neutron capture cross-section (σ), is measured in barns ($1 \text{ b} = 10^{-24} \text{ cm}^2$). Of the
20 various nuclides that have high neutron capture cross-sections, ^{10}B is the most attractive for the following reasons: (a) it is nonradioactive and readily available, comprising approximately 20% of naturally occurring boron: (b) the particles emitted by the capture reaction [$^{10}\text{B}(\text{n},\alpha)^7\text{Li}$] are largely high LET: (c) their path lengths are approximately 1 cell diameter (10-14 μm), theoretically
25 limiting the radiation effect to those tumor cells that have taken up a sufficient amount of ^{10}B and simultaneously sparing normal cells and (d) the extensive chemistry of boron is such that it can be incorporated into a multitude of different chemical structures.

^7Li and α -particles are the primary fission product of the neutron
30 capture reaction with ^{10}B . α -Particles are relatively slow and give rise to closely spaced ionizing events that consist of tracks of sharply defined columns. They have a path length of approximately 10 μm , are high LET, and destroy a wide variety of biologically active molecules including DNA, RNA, and proteins. For

these reasons there is little, if any, cellular repair from α -particle-induced radiation injury.

Since the $^{10}\text{B}(\text{n},\alpha)^7\text{Li}$ reaction will produce a significant biological effect only when there is a sufficient fluence of thermal neutrons and a critical amount of ^{10}B localized around, on, or within the cell, the radiation produced can be extremely localized thereby sparing normal tissue components. Thus, selectivity is simultaneously one of the advantages and disadvantages of BNCT, since it requires delivery of boron-10 to tumor cells in greater amounts than normal cells.

Ideally, boron compounds to be used for BNCT should have a high specificity for malignant cells with concomitantly low concentrations in adjacent normal tissues and blood. Since it is desirable to confine the radiation solely to these cells, an intracellular and optimally intranuclear localization of boron would be preferred.

Several boron-containing derivatives of chlorpromazine have been synthesized (Nakagawa, T. et al., Chem.-Pharm. Bull. (Tokyo), 24, 778-781 (1976); Alam, F. et al., Sthralenther. Onkol., 165, 121-125 (1989)) and are being evaluated for their *in vivo* tumor localizing properties. *p*-Boronophenylalanine is another compound that is being studied as a potential capture agent for the treatment of melanoma. The rationale for its use is the avidity of melanomas for aromatic amino acids and their subsequent incorporation into melanin (Ichihashi, M. et al., J. Invest. Dermatol., 78, 215-218 (1982); Mishima, Y. et al., Neutral Capture Therapy, 230-236, Niigata, Japan: Nishimura Co., Ltd. (1986)).

Tumor localization has been demonstrated following i.v. administration by means of whole body autoradiography (Coderre, J.A. et al., Cancer Res., 48, 6313-6316 (1988)) and in several patients with cutaneous melanoma following perilesional injection (Mishima, Y. et al., Sthralenther. Onkol., 165, 251-254 (1989)). Stimulated by Mishima's experience, a number of other boron-containing amino acids have been synthesized that potentially could be incorporated in larger amounts into proteins of malignant cells (Hall, I.H. et al., J. Pharm. Sci., 68, 685-688 (1979)).

Another approach to the selective targeting of boron to melanomas is based on the observation that thiouracil is preferentially incorporated into melanotic melanomas during melanogenesis (Whittacker, J.R., I. Biol. Chem., **246**, 6217-6226 (1971)). This observation provided the impetus
5 for the synthesis of several boron-containing thiouracils (Gabel, D., Clinical Aspects of Neutron Capture Therapy, 233-241, New York: Plenum Publishing Co. (1989)), and these currently are being evaluated in animals.

Two other classes of compounds with a propensity for localizing in malignant tumors are the porphyrins and the related phthalocyanines. The
10 biochemical basis by which these compounds achieve elevated concentration in malignant tumors is unknown, but this observation has served as the rationale for the use of hematoporphyrin derivative in the photodynamic therapy of cancer (Dougherty, T.J. et al., Porphyrin Photosensitization, 3-13, New York: Plenum Publishing Corp. (1981)).

15 The high concentration of these compounds in tumors and their intracellular localization and persistence have stimulated several groups of investigators to synthesize boronated porphyrins (Kahl, S. B. et al., Neutron Capture Therapy, 61-67, Niigata, Japan: Nishimura Co., Ltd. (1986)) and phthalocyanines (Alam, F. et al., Strahlenther. Oncol., **165**, 121-123 (1989)) as
20 potential capture agents. Boronated porphyrins appear to be 3-4 times more effective per unit dose in cell culture than the monomeric or dimeric form of $\text{Na}_2\text{B}_{12}\text{H}_{11}\text{SH}$ (Laster, B.H. et al., Strahlenther. Oncol., **165**, 203-205 (1989)). Although liver concentrations of these compounds are also high (Kahl, S.B. et al., *supra*) this would not limit their use as a capture agent for the treatment of
25 brain tumors.

One final category of low molecular weight boron compounds are boron-containing purines and pyrimidines and their nucleosides. The rationale for their development is that such compounds may be selectively incorporated into rapidly proliferating tumor cells and trapped within the cell following their
30 conversion to the corresponding nucleotide. Alternatively, these bases and their nucleosides may function as analogues of naturally occurring precursors of nucleic acids and become incorporated into nuclear DNA.

Cytoplasmic or preferably a nuclear localization of all of these boron compounds would be advantageous since the heavy particles resulting from the capture reaction would deliver a greater proportion of their energy to intranuclear targets, thereby permitting lower boron concentrations than would have been required if the compounds were located extracellularly (Gabel, D. et al., Radiat. Res., 111, 14-25 (1987); Fairchild, R.G. et al., Int. J. Radiat. Oncol. Biol. Phys., 11, 831-840 (1985)). Schinazi and Prusoff (Schinazi, R.F. et al., Tetrahedron Lett., 50, 4981-4984 (1978)) have synthesized the first boron-containing nucleoside, 5-dihydroloxyboryl-2'-deoxyuridine, an analogue of thymidine, and have shown that it was not cytotoxic to African green monkey (Vero) cells at a concentration level of 1600 μ M (Laster, B.H. et al., Neutron Capture Therapy, 46-54, Niigata, Japan: Nishimura Co., Ltd. (1986)). *In vitro* neutron radiation studies of cells grown in the presence of 5-dihydroxy-2'-deoxyuridine produced a biological effect that was equivalent to a concentration of $\sim 6 \mu\text{g } ^{10}\text{B/g}$, which, if attainable *in vivo*, would be sufficient for BNCT.

During the 1960s and early 1970s, interests developed in the potential use of polyclonal antibodies directed against tumor-associated antigens for the delivery of drugs and radioisotopes to tumors (Pressman, D. et al., Cancer Res., 40, 3001-3007 (1957); Ghose, T. et al., Br. Med. J., 1, 90-93 (1967); Ghose, T. et al., Cancer (Phila.), 29, 1398-1400 (1972)). In 1964, Soloway suggested that antibodies might be used for the selective targeting of ^{10}B to tumors (Soloway, A.H., *supra*). Hawthorne et al. (Hawthorne, M.F. et al., J. Medicinal Chem., 15, 449-452 (1972)) reported on the incorporation of the diazonium salt from 1-(4-aminophenyl)-1,2-dicarbo-*closo*-dodecaborane into antibodies directed against bovine serum albumin and polyclonal antibodies directed against human and mouse histocompatibility antigens (Hawthorne, M.F. et al., *supra*).

It was claimed from *in vitro* experiments that these immunoconjugates were capable of delivering enough boron to human and murine lymphocytes to sustain a lethal $^{10}\text{B}(\text{n},\alpha)^7\text{Li}$ reaction, as evidenced by reduced viability following neutron irradiation. However, the immunoconjugates contained only 0.2% natural boron by weight, which was equal to 6 atoms of ^{10}B /molecule of antibody. In retrospect, it appears that there

must have been some other explanation for the reduced cell viability that was observed. Sneath et al. (Sneath, R.L., Jr., J. Medicinal Chem., 17, 796-799 (1974)) showed that water-solubilizing groups had to be incorporated into protein-binding polyhedral boranes if protein solubility in aqueous systems was to be maintained.

Subsequently, a group of polyhedral borane derivatives containing protein-binding functional groups were linked to IgG molecules by means of the carbodiimide reaction without evidence of precipitation (Sneath, R.L. et al., J. Medicinal Chem., 19, 1290-1294 (1976)).

One final category of macromolecular species that are possibly useful for the delivery of ^{10}B is what may be termed "encapsulating complexes," such as liposomes, microspheres, and low density lipoproteins (Kahl, S.B. et al., Strahlenther. Onkol., 165, 137-139 (1989)). Theoretically, large amounts of ^{10}B could be encapsulated, and if these encapsulating complexes could be targeted to the tumor by linkage to a monoclonal antibody using existing methodology or targeting an endogenously expressed cell surface receptor, they might be powerful delivery systems. Again, there may be preferential localization in the reticuloendothelial system, and methodology would have to be developed to minimize this and maximize tumor uptake.

Cobalamin

For several years after the isolation of vitamin B_{12} as cyanocobalamin in 1948, it was assumed that cyanocobalamin and possibly hydroxocobalamin, its photolytic breakdown product, occurred in man. Since then it has been recognized that cyanocobalamin is an artifact of the isolation of vitamin B_{12} and that hydroxocobalamin and the two coenzyme forms, methylcobalamin and adenosylcobalamin, are the naturally occurring forms of the vitamin.

The structure of these various forms is shown in Figure 1, wherein X is CN, OH, CH_3 or adenosyl, respectively. Hereinafter, the term cobalamin will be used to refer to all of the molecule except the X group. The fundamental ring system without cobalt (Co) or side chains is called *corrin* and the octadehydrocorrin is called *corrole*. Figure 1 is adapted from The Merck Index, Merck & Co. (11th ed. 1989), wherein X is above the plane defined by the

corrin ring and nucleotide is below the plane of the ring. The corrin ring has attached six amidoalkyl ($\text{H}_2\text{NC(O)Alk}$) substituents, at the 2, 3, 7, 8, 13, and 18 positions, which can be designated a-e and g, respectively. See D.L. Anton et al., J. Amer. Chem. Soc., **102**, 2215 (1980).

5 Methylcobalamin serves as the cytoplasmic coenzyme for ^5N -methyltetrahydrofolate:homocysteine methyl transferase (methionine synthetase, EC 2.1.1.13), which catalyzes the formation of methionine from homocysteine. Adenosylcobalamin is the mitochondrial coenzyme for methylmalonyl CoA mutase (EC 5.4.99.2) which interconverts methylmalonyl CoA and succinyl
10 CoA.

 All forms of vitamin B_{12} (adenosyl-, cyano-, hydroxo-, or methylcobalamin) must be bound by the transport proteins, Intrinsic Factor and Transcobalamin II to be biologically active. Specifically, gastrointestinal absorption of vitamin B_{12} relies upon the intrinsic factor-vitamin B_{12} complex
15 being bound by the intrinsic factor receptors in the terminal ileum. Likewise, intravascular transport and subsequent cellular uptake of vitamin B_{12} throughout the body is dependent upon transcobalamin II and the cell membrane transcobalamin II receptors, respectively. After the transcobalamin II-vitamin B_{12} complex has been internalized, the transport protein undergoes lysozymal
20 degradation, which releases vitamin B_{12} into the cytoplasm. All forms of vitamin B_{12} can then be interconverted into adenosyl-, hydroxo-, or methylcobalamin depending upon cellular demand. See, for example, A.E. Finkler et al., Arch. Biochem. Biophys., **120**, 79 (1967); C. Hall et al., J. Cell Physiol., **133**, 187 (1987); M.E. Rappazzo et al., J. Clin. Invest., **51**, 1915 (1972) and R. Soda et al.,
25 Blood, **65**, 795 (1985).

 Cells undergoing rapid proliferation have been shown to have increased uptake of thymidine and methionine. (See, for example, M.E. van Eijkeren et al., Acta Oncologica, **31**, 539 (1992); K. Kobota et al., J. Nucl. Med., **32**, 2118 (1991) and K. Higashi et al., J. Nucl. Med., **34**, 773 (1993)). Since
30 methylcobalamin is directly involved with methionine synthesis and indirectly involved in the synthesis of thymidylate and DNA, it is not surprising that methylcobalamin as well as Cobalt-57-cyanocobalamin have also been shown to have increased uptake in rapidly dividing tissue (for example, see, B.A. Cooper

et al., Nature, **191**, 393 (1961); H. Flodh, Acta Radiol. Suppl., **284**, 55 (1968); L. Bloomquist et al., Experientia, **25**, 294 (1969)). Additionally, upregulation in the number of transcobalamin II receptors has been demonstrated in several malignant cell lines during their accelerated thymidine incorporation and DNA
5 synthesis (see, J. Lindemans et al., Exp. Cell. Res., **184**, 449 (1989); T. Amagasaki et al., Blood, **26**, 138 (1990) and J.A. Begly et al., J. Cell Physiol., **156**, 43 (1993).

Vitamin B₁₂ has several characteristics which potentially make it an attractive *in vivo* tumor therapeutic agent. Vitamin B₁₂ is water soluble, has
10 no known toxicity, and in excess is excreted by glomerular filtration. In addition, the uptake of vitamin B₁₂ could potentially be manipulated by the administration of nitrous oxide and other pharmacological agents (D. Swanson et al., Pharmaceuticals in Medical Imaging, MacMillan Pub. Co., NY (1990) at pages 621-628).

15 A process for preparing ¹²⁵I-vitamin B₁₂ derivatives is described in U.S. Pat. No. 3,981,863 issued to Niswender et al. In this process, vitamin B₁₂ is first subjected to mild hydrolysis to form a mixture of monocarboxylic acids, which Houts, *infra*, disclosed to contain mostly the (e)-isomer. The mixture is then reacted with a p-(aminoalkyl)phenol to introduce a phenol group into the
20 B₁₂ acids (via reaction with one of the free carboxylic acid groups). The mixed substituent B₁₂ derivatives are then iodinated in the phenol-group substituent. This U.S. patent teaches that the mixed ¹²⁵I-B₁₂ derivatives so made are useful in the radioimmunoassay of B₁₂, using antibodies raised against the mixture.

U.S. Pat. No. 4,465,775 issued to T. M. Houts reported that the
25 components of the radiolabelled mixture of Niswender et al. did not bind with equal affinity to IF. Houts disclosed that radioiodinated derivatives of the pure monocarboxylic (d)-isomer are useful in assays of B₁₂ in which IF is used. However, although Houts generally discloses that the monocarboxylic (d)-isomer can be labelled with fluorophores or enzymes and used in competitive
30 assays for vitamin B₁₂ in fluids, a continuing need exists for labelled vitamin B₁₂ derivatives suitable for tumor and organ imaging and therapy.

U.S. Patent Number 5,739,313 issued to Collins and Hogenkamp reported that cobalamin analogs comprising a linking group and a chelating

group optionally comprising a detectable radionuclide or a paramagnetic ion
localize in tumor cells and are useful for imaging tumors.

Despite previous efforts to identify a neutron capture agent that
localizes in tumor cells in high concentration and is useful to treat cancer, there is
5 currently a need for neutron capture agents that are useful to treat tumors.

Summary of the Invention

The invention provides a compound of the invention which is a
residue of a compound of formula I (Figure 1) I linked to a residue of a molecule
comprising Boron-10 (i.e., B-10), wherein X is CN, OH, CH₃, adenosyl, or a
10 molecule comprising B-10; or a pharmaceutically acceptable salt thereof.

The invention also provides a compound wherein a residue of a
compound of formula I (Figure 1) is linked to a group of the formula Q-L-W-
Det; wherein X is CN, OH, CH₃, adenosyl, a molecule comprising B-10, or Q-L-
W-Det; wherein Det is a chelating group comprising Gd-157; L is a linker or
15 absent; and W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-,
-OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)₂-, -C(=O)-, -N(R)-, or a direct
bond; wherein each R is independently H or (C₁-C₆)alkyl; or a pharmaceutically
acceptable salt thereof.

The invention also provides a compound wherein a residue of a
20 compound of formula I (Figure 1) is linked to a residue of a molecule comprising
B-10; wherein a residue of the compound of formula I is also linked to a group
of the formula Q-L-W-Det, wherein X is CN, OH, CH₃, adenosyl, a group of the
formula Q-L-W-Det, or a molecule comprising B-10; wherein: Det is a chelating
group comprising a therapeutic radionuclide or a diagnostic radionuclide; L is a
25 linker or absent; and Q and W are each independently -N(R)C(=O)-,
-C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)₂-, -C(=O)-, -N(R)-,
or a direct bond; wherein each R is independently H or (C₁-C₆)alkyl; or a
pharmaceutically acceptable salt thereof.

The invention also provides a compound wherein a residue of a
30 compound of formula I (Figure 1) is linked to a residue of a molecule comprising
B-10; wherein the residue of the compound of formula I is also linked to a group
of the formula Q-L-W-Det, wherein X is CN, OH, CH₃, adenosyl, a group of the
formula Q-L-W-Det, or a molecule comprising B-10; Det is a chelating group

comprising Gd-157; L is a linker or absent; and Q and W are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)₂-, -C(=O)-, -N(R)-, or a direct bond; wherein each R is independently H or (C₁-C₆)alkyl; or a pharmaceutically acceptable salt thereof.

5 The invention also provides a compound wherein a residue of a compound of formula I (Figure 1) is linked to a detectable radionuclide and also linked to a residue of a molecule comprising Boron-10 (i.e., B-10), wherein X is CN, OH, CH₃, adenosyl, or a molecule comprising B-10; or a pharmaceutically acceptable salt thereof.

10 The invention also provides a compound wherein a residue of a compound of formula I (Figure 1) is linked to a detectable radionuclide; and is also linked to a group comprising Gd-157; wherein X is CN, OH, CH₃, adenosyl, a molecule comprising B-10, or a group comprising Gd-157; or a pharmaceutically acceptable salt thereof. Specifically, the group comprising Gd-157 can have the formula Q-L-W-Det; wherein Det is a chelating group
15 comprising Gd-157; L is a linker or absent; and W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)₂-, -C(=O)-, -N(R)-, or a direct bond; wherein each R is independently H or (C₁-C₆)alkyl.

20 The invention also provides a compound of wherein a residue of a compound of formula I (Figure 1) is linked 1) to a molecule comprising B-10 or to a chelating group comprising Gd-157, and is linked 2) to at least one residue of the formula Q-L-W-Det; wherein each Det is independently a chelating group comprising a metallic radionuclide; each L is independently a linker or
25 absent; and each W and Q is independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)₂-, -C(=O)-, -N(R)-, or a direct bond; wherein each R is independently H or (C₁-C₆)alkyl; or a pharmaceutically acceptable salt thereof.

30 The invention also provides a pharmaceutical composition comprising a compound of the present invention and a pharmaceutically acceptable carrier.

 The invention also provides a method of treating a tumor in a mammal in need of such treatment comprising administering to the mammal an

effective amount of a compound of the present invention; and administering neutron capture therapy.

The invention also provides a method for imaging a tumor in a mammal in need of such imaging comprising administering to the mammal a
5 detectable amount of a compound of the present invention in combination with a pharmaceutically acceptable vehicle effective to image the tumor; and detecting the presence of the compound.

The invention also provides a compound of the present invention for use in medical therapy or diagnosis.

10 The invention also provides the use of a compound of the present invention for the manufacture of a medicament for imaging a tumor in a mammal (e.g., a human).

The invention also provides the use of a compound of the present invention for the manufacture of a medicament for treating a tumor in a mammal
15 (e.g., a human).

The invention also provides intermediates disclosed herein that are useful in the preparation of the compounds of the present invention as well as synthetic methods useful for preparing the compounds of the invention.

Brief Description of the Figures

20 Figure 1 depicts the structure of cobalamin wherein X is CN (cyano), OH, CH₃ or adenosyl.

Figure 2 illustrates the synthesis of a cyanocobalamin-nido-carborane conjugate.

Figure 3 illustrates the synthesis of representative compounds of
25 the present invention.

Detailed Description of the Invention

Applicant has discovered certain cobalamin analogs are useful, in combination with neutron capture therapy, to treat tumors. Applicant has also discovered certain cobalamin analogs are useful as neutron capture agents and as
30 tumor imaging agents.

The following definitions are used, unless otherwise described: halo is fluoro, chloro, bromo, or iodo. Alkyl, alkoxy, alkenyl, alkynyl, etc. denote both straight and branched groups; but reference to an individual radical

such as "propyl" embraces only the straight chain radical, a branched chain isomer such as "isopropyl" being specifically referred to. Aryl denotes a phenyl radical or an ortho-fused bicyclic carbocyclic radical having about nine to ten ring atoms in which at least one ring is aromatic.

5 Specific and preferred values listed below for radicals, substituents, and ranges, are for illustration only; they do not exclude other defined values or other values within defined ranges for the radicals and substituents.

 It is appreciated that those skilled in the art will recognize that
10 compounds of the present invention having a chiral center may exist in and be isolated in optically active and racemic forms. Some compounds may exhibit polymorphism. It is to be understood that the present invention encompasses any racemic, optically-active, polymorphic, or stereoisomeric form, or mixtures thereof, of a compound of the invention, which possess the useful properties
15 described herein, it being well known in the art how to prepare optically active forms (for example, by resolution of the racemic form by recrystallization techniques, by synthesis from optically-active starting materials, by chiral synthesis, or by chromatographic separation using a chiral stationary phase) and how to determine antitumor activity or utility as an antitumor imaging agent
20 using the standard tests described herein, or using other similar tests which are well known in the art.

 Specifically, (C₁-C₁₄)alkyl can be methyl, ethyl, propyl, isopropyl, butyl, iso-butyl, sec-butyl, pentyl, 3-pentyl, hexyl, heptyl, octyl, nonyl, decyl undecyl, dodecyl, tridecyl or tetradecyl.

25 Specifically, (C₂-C₁₄)alkenyl can be vinyl, allyl, 1-propenyl, 2-propenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 1-heptenyl, 2-heptenyl, 3-heptenyl, 4-heptenyl, 5-heptenyl, 6-heptenyl, 1-octenyl, 2-octenyl, 3-octenyl, 4-octenyl, 5-octenyl, 6-octenyl, 7-octenyl, 1-nonenyl, 2-nonenyl, 3-nonenyl, 4-nonenyl, 5-nonenyl, 6-nonenyl, 7-nonenyl, 8-nonenyl, 1-decenyl, 2-decenyl, 3-decenyl, 4-decenyl, 5-decenyl, 6-decenyl, 7-decenyl, 8-decenyl, 9-decenyl, 1-undecenyl, 2-undecenyl, 3-undecenyl, 4-undecenyl, 5-undecenyl, 6-undecenyl, 7-undecenyl, 8-undecenyl, 9-undecenyl, 10-undecenyl, 1-dodecenyl,
30

2 -dodecenyl, 3 -dodecenyl, 4 -dodecenyl, 5-dodecenyl, 6-dodecenyl, 7-dodecenyl, 8-dodecenyl, 9-dodecenyl, 10-dodecenyl, 11-dodecenyl, 1-tridecenyl, 2-tridecenyl, 3-tridecenyl, 4-tridecenyl, 5-tridecenyl, 6-tridecenyl, 7-tridecenyl, 8-tridecenyl, 9-tridecenyl, 10-tridecenyl, 11-tridecenyl, 12-tridecenyl, 1-tetradecenyl, 2-tetradecenyl, 3-tetradecenyl, 4-tetradecenyl, 5-tetradecenyl, 6-tetradecenyl, 7-tetradecenyl, 8-tetradecenyl, 9-tetradecenyl, 10-tetradecenyl, 11-tetradecenyl, 12-tetradecenyl or 13-tetradecenyl.

Specifically, (C₂-C₁₄)alkynyl can be ethynyl, 1-propynyl, 2-propynyl, 1-butylnyl, 2-butylnyl, 3-butylnyl, 1-pentylnyl, 2-pentylnyl, 3-pentylnyl, 4-pentylnyl, 1-hexynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 5-hexynyl, 1-heptylnyl, 2-heptylnyl, 3-heptylnyl, 4-heptylnyl, 5-heptylnyl, 6-heptylnyl, 1-octynyl, 2-octynyl, 3-octynyl, 4-octynyl, 5-octynyl, 6-octynyl, 7-octynyl, 1-nonylnyl, 2-nonylnyl, 3-nonylnyl, 4-nonylnyl, 5-nonylnyl, 6-nonylnyl, 7-nonylnyl, 8-nonylnyl, 1-decynyl, 2-decynyl, 3-decynyl, 4-decynyl, 5-decynyl, 6-decynyl, 7-decynyl, 8-decynyl, 9-decynyl, 1-undecynyl, 2-undecynyl, 3-undecynyl, 4-undecynyl, 5-undecynyl, 6-undecynyl, 7-undecynyl, 8-undecynyl, 9-undecynyl, 10-undecynyl, 11-undecynyl, 1-dodecynyl, 2 -dodecynyl, 3 -dodecynyl, 4 -dodecynyl, 5-dodecynyl, 6-dodecynyl, 7-dodecynyl, 8-dodecynyl, 9-dodecynyl, 10-dodecynyl, 11-dodecynyl, 1-tridecynyl, 2-tridecynyl, 3-tridecynyl, 4-tridecynyl, 5-tridecynyl, 6-tridecynyl, 7-tridecynyl, 8-tridecynyl, 9-tridecynyl, 10-tridecynyl, 11-tridecynyl, 12-tridecynyl, 1-tetradecynyl, 2-tetradecynyl, 3-tetradecynyl, 4-tetradecynyl, 5-tetradecynyl, 6-tetradecynyl, 7-tetradecynyl, 8-tetradecynyl, 9-tetradecynyl, 10-tetradecynyl, 11-tetradecynyl, 12-tetradecynyl or 13-tetradecynyl.

Specifically "aryl" can be phenyl, indenyl, or naphthyl.

Specifically (C₃-C₈)cycloalkyl can be cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl.

As used herein, an "amino acid" is a natural amino acid residue (e.g. Ala, Arg, Asn, Asp, Cys, Glu, Gln, Gly, His, Hyl, Hyp, Ile, Leu, Lys, Met, Phe, Pro, Ser, Thr, Trp, Tyr, and Val) in D or L form, as well as unnatural amino acid (e.g. phosphoserine; phosphothreonine; phosphotyrosine; hydroxyproline; gamma-carboxyglutamate; hippuric acid; octahydroindole-2-carboxylic acid; statine; 1,2,3,4,-tetrahydroisoquinoline-3-carboxylic acid; penicillamine; ornithine; citruline; α-methyl-alanine; para-benzoylphenylalanine;

phenylglycine; propargylglycine; sarcosine; and tert-butylglycine) residue having one or more open valences. The term also comprises natural and unnatural amino acids bearing conventional amino protecting groups (e.g. acetyl, acyl, trifluoroacetyl, or benzyloxycarbonyl), as well as natural and unnatural amino acids protected at carboxy (e.g. as a (C₁-C₆)alkyl, phenyl or benzyl ester or amide) with conventional protecting groups. Other suitable amino and carboxy protecting groups are known to those skilled in the art (See for example, T.W. Greene, Protecting Groups In Organic Synthesis; Wiley: New York, 1981; D. Voet, Biochemistry, Wiley: New York, 1990; L. Stryer, Biochemistry, (3rd Ed.), W.H. Freeman and Co.: New York, 1975; J. March, Advanced Organic Chemistry, Reactions, Mechanisms and Structure, (2nd Ed.), McGraw Hill: New York, 1977; F. Carey and R. Sundberg, Advanced Organic Chemistry, Part B: Reactions and Synthesis, (2nd Ed.), Plenum: New York, 1977; and references cited therein). According to the invention, the amino or carboxy protecting group can also comprise a radionuclide (e.g., Fluorine-18, Iodine-123, or Iodine-124).

As used herein, a "peptide" is a sequence of 2 to 25 amino acids (e.g. as defined hereinabove) or peptidic residues having one or more open valences. The sequence may be linear or cyclic. For example, a cyclic peptide can be prepared or may result from the formation of disulfide bridges between two cysteine residues in a sequence. A peptide can be linked through the carboxy terminus, the amino terminus, or through any other convenient point of attachment, such as, for example, through the sulfur of a cysteine. Peptide derivatives can be prepared as disclosed in U.S. Patent Numbers 4,612,302; 4,853,371; and 4,684,620, or as described in the Examples hereinbelow. Peptide sequences specifically recited herein are written with the amino terminus on the left and the carboxy terminus on the right.

Particular and specific values listed below for radicals, substituents, and ranges, are for illustration only and they do not exclude other defined values or other values within defined ranges for the radicals and substituents.

Specifically, the peptide can be poly-L-lysine, poly-L-glutamic acid, poly-L-aspartic acid, poly-L-histidine, poly-L-ornithine, poly-L-serine,

poly-L-threonine, poly-L-tyrosine, poly-L-lysine--L-phenylalanine or poly-L-lysine-L-tyrosine.

As used herein, a "residue of a compound of formula I" is a radical of a compound of formula I having one or more open valences. Any synthetically feasible atom or atoms of the compound of formula I may be removed to provide the open valence, provided the resulting compound is able to localize in or near tumors. Based on the linkage that is desired, one skilled in the art can select suitably functionalized starting materials that can be derived from a compound of formula I using procedures that are known in the art. For example, suitable atoms that may be removed include the NH₂ group of the a-carboxamide (illustrated in figure 1) or a hydrogen atom from the NH₂ group of the a-carboxamide, the NH₂ group of the b-carboxamide (illustrated in figure 1) or a hydrogen atom from the NH₂ group of the b-carboxamide, the NH₂ group of the d-carboxamide (illustrated in figure 1) or a hydrogen atom from the NH₂ group of the d-carboxamide, the NH₂ group of the e-carboxamide (illustrated in figure 1) or a hydrogen atom from the NH₂ group of the e-carboxamide, and X at the 6-position (illustrated in figure 1). In addition, the hydrogen atom of the hydroxy group at the 3' position of the sugar, the hydrogen atom from the hydroxyl group at the 3' position of the sugar, the hydrogen atom of the CH₂OH group at the 5' position, or the hydrogen atom from the hydroxyl group at the 5' position of the sugar ring may be removed.

As used herein, "adenosyl" is an adenosine radical in which any synthetically feasible atom or group of atoms have been removed, thereby providing an open valence. Synthetically feasible atoms which may be removed include the hydrogen atom of the hydroxy group at the 5' position. Accordingly, adenosyl can conveniently be attached to the 6-position of a compound of formula I via the 5' position of adenosyl.

As used herein, a "molecule comprising B-10" can be any compound that contains at least one B-10 atom. The nature of the molecule that includes B-10 is not critical. The compound, however, must be nontoxic and must be able to enter the tumor cell or locate near the tumor cell when the molecule comprising B-10 is attached.

As used herein, a "residue of a molecule comprising B-10" is a radical of a molecule comprising B-10 having one or more open valences. Any synthetically feasible atom or atoms of the molecule comprising B-10 may be removed to provide the open valence, provided bioactivity is substantially retained. Based on the linkage that is desired, one skilled in the art can select suitably functionalized starting materials that can be derived from a molecule comprising B-10 using procedures that are known in the art.

As used herein, a "residue of o-carborane," a "residue of m-carborane," or a "residue of p-carborane" is a radical of o-carborane, m-carborane or p-carborane, respectively, having one or more open valences. Any synthetically feasible atom or atoms of o-carborane, m-carborane or p-carborane may be removed to provide the open valence, provided bioactivity is substantially retained. Based on the linkage that is desired, one skilled in the art can select suitably functionalized starting materials that can be derived from o-carborane, m-carborane or p-carborane using procedures that are known in the art.

Boron Compounds Useful For Neutron Capture Therapy

The invention provides a compound of formula I (Figure 1) linked to one or more molecules comprising B-10, wherein X is CN, OH, CH₃, adenosyl or a molecule comprising B-10; or a pharmaceutically acceptable salt thereof.

A variety of molecules comprising B-10 known in the art are useful in the present invention. The molecules vary considerably in structure but are suitable to practice the present invention. Acceptable species include boron containing amino acids, carbohydrates, and nucleosides, as well as carboranes. A wide variety of boron-containing compounds are commercially available, or are known in the art. A variety of molecules comprising B-10 are commercially available from Boron Biologicals, Inc., Raleigh, North Carolina and RysCor Science, Inc., Raleigh, North Carolina.

Specifically, at least one molecule comprising B-10 can be o-carborane, m-carborane or p-carborane. More specifically, at least one molecule comprising B-10 is o-carborane. o-Carborane [1,2-dicarbadoecaborane(12)]; m-carborane [1,7-dicarbadoecaborane(12)]; and p-carborane [1,12-

dicarbadodecaborane(12)] are commercially available from Aldrich, Milwaukee, WI.

Specifically, each molecule comprising B-10 can independently be o-carborane, m-carborane or p-carborane. More specifically, each molecule
5 comprising B-10 is o-carborane.

Compound of Formula I / Molecule Comprising B-10

The residue of molecule comprising B-10 can be linked to the residue of a compound of formula I through an amide (e.g., -N(R)C(=O)- or -C(=O)N(R)-), ester (e.g., -OC(=O)- or -C(=O)O-), ether (e.g., -O-), amino (e.g.,
10 -N(R)-), ketone (e.g., -C(=O)-), thioether (e.g., -S-), sulfinyl (e.g., -S(O)-), sulfonyl (e.g., -S(O)₂-), or a direct (e.g., C-C bond) linkage, wherein each R is independently H or (C₁-C₆)alkyl. Such a linkage can be formed from suitably functionalised starting materials using synthetic procedures that are known in the art. Based on the linkage that is desired, one skilled in the art can select suitably
15 functional starting materials that can be derived from a residue of a compound of formula I and from a given residue of a molecule comprising B-10 using procedures that are known in the art.

The residue of the molecule comprising B-10 can be directly linked to any synthetically feasible position on the residue of a compound of
20 formula I. Suitable points of attachment include, for example, the b-carboxamide, the d-carboxamide, and the e-carboxamide (illustrated in figure 1), as well as the 6-position (the position occupied by X in figure 1), and the 5'-hydroxy and the 3'-hydroxy groups on the 5-membered sugar ring, although other points of attachment are possible. U.S. Patent No. 5,739,313 discloses
25 compounds (e.g., cyanocobalamin-b-(4-aminobutyl)amide, methylcobalamin-b-(4-aminobutyl)amide, and adenosylcobalamin-b-(4-aminobutyl)amide) that are useful intermediates for the preparation of compounds of the present invention.

Compounds wherein the residue of a molecule comprising B-10 is linked to the 6-position of a compound of formula I can be prepared by reducing
30 a corresponding Co (II) compound of formula I to form a nucleophilic Co (I) compound and treating this Co (I) compound with a residue of a molecule comprising B-10 (or a derivative thereof) comprising a suitable leaving group, such as a halide (e.g., a chloride).

The invention also provides compounds having more than one residue of a molecule comprising B-10 directly linked to a compound of formula I. For example, the residue of a molecule comprising B-10 can be directly linked to a residue of the b-carboxamide of the compound of formula I and a residue of another molecule comprising B-10 can be directly linked to a residue of the d-carboxamide of the compound of formula I. In addition, the residue of a molecule comprising B-10 can be directly linked to the 6-position of the compound of formula I and a residue of another molecule comprising B-10 can be directly linked to a residue of the d- or e-carboxamide of the compound of formula I.

Compound of Formula I / Linker / Molecule Comprising B-10 Linker

In addition to being directly linked to the residue of a compound of formula I, the residue of a molecule comprising B-10 can also be linked to the residue of a compound of formula I by a suitable linker. The structure of the linker is not crucial, provided it yields a compound of the invention which has an effective therapeutic index against the target cells, and which will localize in or near tumor molecules.

Suitable linkers include linkers that separate the residue of a compound of formula I and the residue of a molecule comprising B-10 by about 5 angstroms to about 200 angstroms. Other suitable linkers include linkers that separate the residue of a compound of formula I and the residue of a molecule comprising B-10 by about 5 angstroms to about 100 angstroms, as well as linkers that separate the residue of a compound of formula I and the residue of a molecule comprising B-10 by about 5 angstroms to about 50 angstroms, or by about 5 angstroms to about 25 angstroms. Suitable linkers are disclosed, for example, in U.S. Patent No. 5,735,313.

The linker can be linked to any synthetically feasible position on the residue of a compound of the residue of formula I. Suitable points of attachment include, for example, a residue of the b-carboxamide, a residue of the d-carboxamide, and a residue of the e-carboxamide, the 6-position (i.e., the position occupied by X in the compound of formula I), as well as a residue of the 5'-hydroxy group and a residue of the 3'-hydroxy group on the 5-membered sugar ring, although other points of attachment are possible. Based on the

linkage that is desired, one skilled in the art can select suitably functionalized starting materials that can be derived from a compound of formula I and from a given molecule comprising B-10 using procedures that are known in the art.

The linker can conveniently be linked to the residue of a compound of formula I or to the residue of a molecule comprising B-10 through an amide (e.g., -N(R)C(=O)- or -C(=O)N(R)-), ester (e.g., -OC(=O)- or -C(=O)O-), ether (e.g., -O-), ketone (e.g., -C(=O)-) thioether (e.g., -S-), sulfinyl (e.g., -S(O)-), sulfonyl (e.g., -S(O)₂-), amino (e.g., -N(R)-) or a direct (e.g., C-C) linkage, wherein each R is independently H or (C₁-C₆)alkyl. The linkage can be formed from suitably functionalised starting materials using synthetic procedures that are known in the art. Based on the linkage that is desired, one skilled in the art can select suitably functional starting materials that can be derived from a residue of a compound of formula I, a residue of a molecule comprising B-10, and from a given linker using procedures that are known in the art.

Specifically, the linker can be a divalent radical of the formula W-A-Q wherein A is (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, (C₃-C₈)cycloalkyl, or (C₆-C₁₀)aryl, wherein W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)₂-, -N(R)-, -C(=O)-, or a direct bond (i.e., W or Q is absent); wherein each R is independently H or (C₁-C₆)alkyl.

Specifically, the linker can be a divalent radical of the formula W-(CH₂)_n-Q wherein, n is between about 1 and about 20, between about 1 and about 15, between about 2 and about 10, between about 2 and about 6, or between about 4 and about 6; wherein W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)₂-, -C(=O)-, -N(R)-, or a direct bond (i.e., W or Q is absent); wherein each R is independently H or (C₁-C₆)alkyl.

Specifically, W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -N(R)-, -C(=O)O-, -O-, or a direct bond (i.e., W or Q is absent).

Specifically, the linker is a divalent radical, i.e., 1,ω-divalent radicals formed from a peptide or an amino acid. The peptide can comprise 2 to about 20 amino acids, 2 to about 15 amino acids, or 2 to about 12 amino acids.

Specifically, the peptide can be poly-L-lysine (i.e., $[-\text{NHCH}[(\text{CH}_2)_4\text{NH}_2]\text{CO}]_m\text{-Q}$, wherein Q is H, $(\text{C}_1\text{-C}_{14})$ alkyl, or a suitable carboxy protecting group; and wherein m is about 2 to about 20. Specifically, the poly-L-lysine can contains about 5 to about 15 residues (i.e., m is between about 5 and about 15). More specifically, the poly-L-lysine can contain about 8 to about 11 residues (i.e., m is between about 8 and about 11).


Specifically, the peptide can be poly-L-glutamic acid, poly-L-aspartic acid, poly-L-histidine, poly-L-ornithine, poly-L-serine, poly-L-threonine, poly-L-tyrosine, poly-L-lysine-L-phenylalanine or poly-L-lysine-L-tyrosine.

Specifically, the linker is prepared from 1,6-diaminohexane $\text{H}_2\text{N}(\text{CH}_2)_6\text{NH}_2$, 1,5-diaminopentane $\text{H}_2\text{N}(\text{CH}_2)_5\text{NH}_2$, 1,4-diaminobutane $\text{H}_2\text{N}(\text{CH}_2)_4\text{NH}_2$, or 1,3-diaminopropane $\text{H}_2\text{N}(\text{CH}_2)_3\text{NH}_2$.

The linker can comprise one or more non-metallic radionuclides. Specifically, the linker can comprise more than one non-metallic radionuclides. More specifically, the linker can comprise 2 to about 10, 2 to about 8, 2 to about 6, or 2 to about 4 non-metallic radioisotopes.

A specific residue of a peptide (i.e., linker) comprising one or more non-metallic radionuclides has the following formula

wherein each M is independently a non-metallic radionuclide; each R is independently $(\text{C}_1\text{-C}_{14})$ alkyl, $(\text{C}_2\text{-C}_{14})$ alkenyl, $(\text{C}_2\text{-C}_{14})$ alkynyl, $(\text{C}_1\text{-C}_{14})$ alkoxy, hydroxy, cyano, nitro, halo, trifluoromethyl, $\text{N}(\text{R}_a)(\text{R}_b)$, $(\text{C}_1\text{-C}_{14})$ alkanoyl, $(\text{C}_2\text{-C}_{14})$ alkanoyloxy, $(\text{C}_6\text{-C}_{10})$ aryl, or $(\text{C}_3\text{-C}_8)$ cycloalkyl wherein R_a and R_b are each independently H or $(\text{C}_1\text{-C}_{14})$ alkyl; P; Q is H, $(\text{C}_1\text{-C}_{14})$ alkyl, or a suitable carboxy protecting group; n is 2 to about 20; i is 1-5, j is 0-4 and $i+j$ is ≤ 5 ; or a pharmaceutically acceptable salt thereof.



Specifically, i can be 1, j can be 0, M can be Fluorine-18, Bromine-76, or Iodine-123, and n can be about 6 to about 12.

The molecule comprising B-10 can comprise one or more boron atoms. A suitable molecule comprising B-10 can contain 1 to about 20, 1 to about 15, 1 to about 10, or 1 to about 5 boron atoms. In addition, the molecule comprising B-10 can comprise one or more B-10 atoms. A suitable molecule
5 comprising B-10 can contain 1 to about 20, 1 to about 15, 1 to about 10, or 1 to about 5 B-10 atoms.

The molecule comprising B-10 can be an amino acid, a carbohydrate, a nucleoside or a carborane. Specifically, the molecule comprising B-10 is o-carborane, m-carborane or p-carborane.

10 Compounds wherein the linker is linked to the 6-position of a compound of formula I can be prepared by preparing a nucleophilic Co (I) species as described herein above, and reacting it with a linker comprising a suitable leaving group, such as a halide (e.g. a chloride).

The invention also provides compounds having more than one
15 molecule comprising B-10 attached to a compound of formula I, each through a linker. For example, the residue of a molecule comprising B-10 can conveniently be linked, through a linker, to a residue of the b-carboxamide of the compound of formula I and a residue of another molecule comprising B-10 can conveniently be linked, through a linker, to a residue of the d- or e-carboxamide
20 of the compound of formula I. In addition, the residue of a molecule comprising B-10 can conveniently be linked, through a linker, to the 6-position of the compound of formula I and a residue of another molecule comprising B-10 can conveniently be linked, through a linker, to a residue of the b-, d- or e-carboxamide of the compound of formula I.

25 The invention also provides compounds having more than one molecule comprising B-10 attached to a compound of formula I, either directly or through a linker. For example, the residue of a molecule comprising B-10 can conveniently be linked, either directly or through a linker, to a residue of the b-carboxamide of the compound of formula I and a residue of another molecule
30 comprising B-10 can conveniently be linked, either directly or through a linker, to a residue of the d- or e-carboxamide of the compound of formula I. In addition, the residue of a molecule comprising B-10 can conveniently be linked, either directly or through a linker, to the 6-position of the compound of formula I

and a residue of another molecule comprising B-10 can conveniently be linked, either directly or through a linker, to a residue of the b-, d- or e-carboxamide of the compound of formula I.

Compound of Formula I / Chelating Group Comprising Gadolinium-157

5 U.S. Patent Number 5,739,313 discloses cobalamin analogs comprising a compound of formula I, a linking group, and a chelating group comprising a detectable radionuclide or a paramagnetic ion. The compounds are disclosed to localize in tumor cells following administration, and to be useful for imaging tumors.

10 The metallic radionuclide Gadolinium-157 is an especially useful ion for conducting magnetic resonance imaging. It is also a useful target ion for neutron capture therapy. Applicants have discovered that incorporation of Gd-157 into one of the chelating molecules disclosed in U.S. Patent Number 5,739,313, provides a compound that is not only particularly useful for
15 conducting magnetic resonance imaging, but also a compound that can be used in conjunction with neutron capture therapy, to treat tumors.

Thus, the present invention provides a residue of a compound of formula I linked to at least one residue of the formula -Q-L-W-Det; wherein each Det is independently a chelating group comprising Gd-157; each L is
20 independently linker (as defined hereinabove) or is absent; and each W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)₂-, -C(=O)-, -N(R)-, or a direct bond (i.e., W or Q is absent); wherein each R is independently H or (C₁-C₆)alkyl.

Compound of Formula I / Chelating Group Comprising a Radionuclide

25 Applicant has also discovered that it is possible to prepare a compound that is useful for both imaging and for treating tumors by incorporating one or more neutron capture target atoms (e.g. B-10 or Gd-157) into a compound that also comprises a detectable radionuclide. Accordingly, the invention provides a residue of a compound of formula I which is linked to one
30 or more residues of a molecule comprising B-10; and which is also linked to one or more groups of the formula -Q-L-W-Det; wherein each Det is independently a chelating group (as defined herein) comprising Gd-157; each L is independently linker (as defined herein) or is absent; and each W and Q are each independently

-N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)₂-, -C(=O)-, -N(R)-, or a direct bond (i.e., W or Q is absent); wherein each R is independently H or (C₁-C₆)alkyl; or a pharmaceutically acceptable salt thereof.

The present invention provides a residue of a compound of
 5 formula I linked to a molecule comprising B-10 or linked to a chelating group comprising Gd-157 and wherein the residue of a compound of formula I is linked to at least one residue of the formula -Q-L-W-Det; wherein each Det is independently a chelating group (as defined hereinabove) comprising a metallic radionuclide; each L is independently a linker (as defined hereinabove) or is
 10 absent; and each W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)₂-, -C(=O)-, -N(R)-, or a direct bond (i.e., W or Q is absent); wherein each R is independently H or (C₁-C₆)alkyl.

Compound of Formula I / Molecule Comprising B-10 / Detectable Radionuclide.

15 The invention also provides a compound wherein a residue of a compound of formula I (Figure 1) is linked 1) to a residue of a molecule comprising B-10, and 2) to a detectable radionuclide, wherein X is CN, OH, CH₃, adenosyl, or a molecule comprising B-10; or a pharmaceutically acceptable salt thereof.

20 The detectable radionuclide can be directly linked to the residue of a compound of formula I, or can be linked by a linker to the residue of a compound of formula I. The linker can be any suitable linker described herein. Specifically, the linker can be of the formula W-A wherein A is (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, (C₃-C₈)cycloalkyl, or (C₆-C₁₀)aryl, wherein W is
 25 -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)₂-, -N(R)-, -C(=O)-, or a direct bond; wherein each R is independently H or (C₁-C₆)alkyl; and wherein A is substituted with one or more non-metallic radionuclides. The linker can also be a peptide or an amino acid.

Compound of Formula I / Group comprising Gd-157 / Detectable Radionuclide.

The invention also provides a compound wherein a residue of a compound of formula I (Figure 1) is linked 1) to a detectable radionuclide; and 2) to a group comprising Gd-157; or a pharmaceutically acceptable salt thereof.

5 The Gd-157 can be linked to the residue of a compound of formula I by any suitable means. For example, the residue of a compound of formula I can be attached to a group comprising Gd-157 that has the formula Q-L-W-Det, wherein Det is a chelating group comprising Gd-157, L is a linker or absent, and W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-, 10 -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)₂-, -C(=O)-, -N(R)-, or a direct bond; wherein each R is independently H or (C₁-C₆)alkyl. Any suitable chelator can be used.

The detectable radionuclide can be directly linked to the residue of a compound of formula I, or can be linked by a linker to the residue of a 15 compound of formula I. The linker can be any suitable linker described herein. Specifically, the linker can be is of the formula W-A wherein A is (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, (C₃-C₈)cycloalkyl, or (C₆-C₁₀)aryl, wherein W is -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)₂-, -N(R)-, -C(=O)-, or a direct bond; wherein each R is independently H or (C₁- 20 C₆)alkyl; and wherein A is substituted with one or more non-metallic radionuclides. The linker can also be a peptide or an amino acid

Non-Metallic Radionuclide

Any detectable non-metallic radionuclide that is suitable for imaging can be used in the compounds of the invention. For example, suitable 25 non-metallic radionuclides include Carbon-11, Fluorine-18, Bromine-76, and Iodine-123. Specifically, the non-metallic radionuclide can be a non-metallic paramagnetic atom (e.g., Fluorine-19); or a non-metallic positron emitting radionuclide (e.g., Carbon-11, Fluorine-18, Iodine-123, or Bromine-76).

Metallic Radionuclide

30 Suitable metallic radionuclides (i.e., metallic radioisotopes or metallic paramagnetic ions) include Antimony-124, Antimony-125, Arsenic-74, Barium-103, Barium-140, Beryllium-7, Bismuth-206, Bismuth-207, Cadmium-109, Cadmium-115m, Calcium-45, Cerium-139, Cerium-141, Cerium-144,

Cesium-137, Chromium-51, Cobalt-55, Cobalt-56, Cobalt-57, Cobalt-58,
 Cobalt-60, Cobalt-64, Copper-67, Erbium-169, Europium-152, Gallium-64,
 Gallium-68, Gadolinium-153, Gadolinium-157 Gold-195, Gold-199, Hafnium-
 175, Hafnium-175-181, Holmium-166, Indium-110, Indium-111, Iridium-192,
 5 Iron-55, Iron-59, Krypton-85, Lead-210, Manganese-54, Mercury-197, Mercury-
 203, Molybdenum-99, Neodymium-147, Neptunium-237, Nickel-63, Niobium-
 95, Osmium-185 + 191, Palladium-103, Platinum-195m, Praseodymium-143,
 Promethium-147, Protactinium-233, Radium-226, Rhenium-186, Rhenium-188,
 Rubidium-86, Ruthenium-103, Ruthenium-106, Scandium-44, Scandium-46,
 10 Selenium-75, Silver-110m, Silver-111, Sodium-22, Strontium-85, Strontium-89,
 Strontium-90, Sulfur-35, Tantalum-182, Technetium-99m, Tellurium-125,
 Tellurium-132, Thallium-204, Thorium-228, Thorium-232, Thallium-170, Tin-
 113, Tin-114, Tin-117m, Titanium-44, Tungsten-185, Vanadium-48, Vanadium-
 49, Ytterbium-169, Yttrium-86, Yttrium-88, Yttrium-90, Yttrium-91, Zinc-65,
 15 and Zirconium-95.


As used herein, a "detectable radionuclide" is any suitable
 radionuclide (i.e., radioisotope) capable of detecting cancer or other neoplastic
 cells in a diagnostic procedure *in vivo* or *in vitro*. Suitable detectable
 radionuclides include metallic radionuclides (i.e., metallic radioisotopes) and
 20 non-metallic radionuclides (i.e., non-metallic radioisotopes).

As used herein, a "therapeutic radionuclide" is any suitable
 radionuclide (i.e., radioisotope) that possesses therapeutic efficacy against cancer
 or other neoplastic cells *in vivo* or *in vitro*. Suitable therapeutic radionuclides
 include metallic radionuclides (i.e., metallic radioisotopes).

25 Chelating groups

Any suitable chelating group can be incorporated into the
 compounds of the invention. Suitable chelating groups include those disclosed
 in U.S. Patent Number 5,739,313. Specifically, the chelating group can be NTA,
 HEDTA, DCTA, RP414, MDP, DOTATOC, CDTA, HYNIC, EDTA, DTPA,
 30 TETA, DOTA, DOTMP, DCTA, 15N4, 9N3, 12N3, or MAG3 (or another
 suitable polyamino acid chelator), which are described herein below, or a
 phosphonate chelator (e.g. EDMT). More specifically, the chelating group can
 be DTPA.

DTPA is diethylenetriaminepentaacetic acid; TETA is 1,4,8,11-tetraazacyclotetradecane-N,N',N'',N'''-tetraacetic acid; DOTA is 1,4,7,10-tetraazacyclododecane-N,N',N'',N'''-tetraacetic acid; 15N4 is 1,4,8,12-tetraazacyclopentadecane-N,N',N'',N'''-tetraacetic acid; 9N3 is 1,4,7-triazacyclononane-N,N',N''-triacetic acid; 12N3 is 1,5,9-triazacyclododecane-N,N',N''-triacetic acid; MAG3 is (N-[N-[N-[(benzoylthio)acetyl]glycyl]glycyl]glycine); and DCTA is a cyclohexane-based metal chelator of the formula



wherein R^3 may be (C_1-C_4) alkyl or $CH_2CO_2^-$, which may be attached through positions 4 or 5, or through the group R^3 and which carries from 1 to 4 detectable metal or nonmetal cations (M), monovalent cations, or the alkaline earth metals. Thus, with metals of oxidation state +1, each individual cyclohexane-based molecule may carry up to 4 metal cations (where both R^3 groups are CH_2COOM). As is more likely, with higher oxidation states, the number of metals will decrease to 2 or even 1 per cyclohexane skeleton. This formula is not intended to limit the molecule to any specific stereochemistry.

NTA, HEDTA, and DCTA are disclosed in Poster Sessions, Proceedings of the 46th Annual Meeting, J. Nuc. Med., p. 316, No. 1386. RP414 is disclosed in Scientific Papers, Proceedings of the 46th Annual Meeting, J. Nuc. Med., p. 123, No. 499. MDP is disclosed in Scientific Papers, Proceedings of the 46th Annual Meeting, J. Nuc. Med., p. 102, No. 413. DOTATOC is disclosed in Scientific Papers, Proceedings of the 46th Annual Meeting, J. Nuc. Med., p. 102, No. 414 and Scientific Papers, Proceedings of the 46th Annual Meeting, J. Nuc. Med., p. 103, No. 415. CDTA is disclosed in Poster Sessions, Proceedings of the 46th Annual Meeting, J. Nuc. Med., p. 318, No.

1396. HYNIC is disclosed in Poster Sessions, Proceedings of the 46th Annual Meeting, J. Nuc. Med., p. 319, No. 1398.

Bifunctional chelators (i.e., chelating groups) based on macrocyclic ligands in which conjugation is via an activated arm attached to the carbon backbone of the ligand can also be employed as a chelating group, as described by M. Moi et al., J. Amer. Chem. Soc., 49, 2639 (1989) (2-p-nitrobenzyl-1,4,7,10-tetraazacyclododecane-N,N',N'',N'''-tetraacetic acid); S. V. Deshpande et al., J. Nucl. Med., 31, 473 (1990); G. Kuser et al., Bioconj. Chem., 1, 345 (1990); C. J. Broan et al., J. C. S. Chem. Comm., 23, 1739 (1990); and C. J. Anderson et al., J. Nucl. Med. 36, 850 (1995) (6-bromoacetamido-benzyl-1,4,8,11-tetraazacyclotetradecane-N,N',N'',N'''-tetraacetic acid (BAT)).

In addition, the diagnostic chelator or therapeutic chelating groups can be any of the chelating groups disclosed in Scientific Papers, Proceedings of the 46th Annual Meeting, J. Nuc. Med., Wednesday, June 9, 1999, p. 124, No. 500.

A "detectable chelating group" is a chelating group comprising a metallic radionuclide (e.g., a metallic radioisotope) capable of detecting cancer or other neoplastic cells *in vivo* or *in vitro*.

Specifically, the chelating group can be any one of the carbonyl complexes disclosed in Waibel et al., Nature Biotechnology, 897-901, Vol. 17, September 1999; or Sattelberger et al., Nature Biotechnology, 849-850, Vol. 17, September 1999.

Specifically, the detectable chelating group can be any of the carbonyl complexes disclosed in Waibel et al., Nature Biotechnology, 897-901, Vol. 17, September 1999; or Sattelberger et al., Nature Biotechnology, 849-850, Vol. 17, September 1999, further comprising a metallic radionuclide. More specifically, the detectable chelating group can be any of the carbonyl complexes disclosed in Waibel et al., Nature Biotechnology, 897-901, Vol. 17, September 1999; or Sattelberger et al., Nature Biotechnology, 849-850, Vol. 17, September 1999, further comprising Technetium-99m.

As used herein, a "therapeutic chelating group" is a chelating group comprising a metallic radionuclide (e.g., a metallic radioisotope) that

possesses therapeutic efficacy against cancer or other neoplastic cells *in vivo* or *in vitro*. Any suitable chelating group can be employed.

Specifically, the therapeutic chelating group can be any of the carbonyl complexes disclosed in Waibel et al., *Nature Biotechnology*, 897-901, Vol. 17, September 1999; or Sattelberger et al., *Nature Biotechnology*, 849-850, Vol. 17, September 1999, further comprising a metallic radionuclide. More specifically, the therapeutic chelating group can be any of the carbonyl complexes disclosed in Waibel et al., *Nature Biotechnology*, 897-901, Vol. 17, September 1999; or Sattelberger et al., *Nature Biotechnology*, 849-850, Vol. 17, September 1999, further comprising Rhenium-186 or Rhenium-188.

Tumors treatable with the compounds and methods of the invention can be located in any part of the mammal. Specifically, the tumor can be located in the breast, lung, thyroid, lymph node, genitourinary system (e.g., kidney, ureter, bladder, ovary, teste, or prostate), musculoskeletal system (e.g., bones, skeletal muscle, or bone marrow), gastrointestinal tract (e.g., stomach, esophagus, small bowel, colon, rectum, pancreas, liver, or smooth muscle), central or peripheral nervous system (e.g., brain, spinal cord, or nerves), head and neck tumors (e.g., ears, eyes, nasopharynx, oropharynx, or salivary glands), or the heart.

The compounds disclosed herein can be prepared using procedures similar to those described in U.S. Patent Number 5,739,313, or using procedures similar to those described herein. The residue of a molecule comprising B-10 can be linked to the residue of a compound of formula I as described hereinabove. Additional intermediates and synthetic procedures useful for preparing compounds of the invention are disclosed, for example, in Hogenkamp, H. et al., *Synthesis and Characterization of nido-Carborane-Cobalamin Conjugates*, Nucl. Med. & Biol., 2000, 27, 89-92; Collins, D., et al., *Tumor Imaging Via Indium 111-Labeled DTPA-Adenosylcobalamin*, Mayo Clinic Proc., 1999, 74:687-691; U.S. Application Ser. No. 60/129,733 filed 16 April 1999; U.S. Application Ser. No. 60/159,874 filed 15 October 1999; U.S. Application Ser. No. 60/159,753 filed 15 October 1999; U.S. Application Ser. No. 60/159,873 filed 15 October 1999; and references cited therein.

In cases where compounds are sufficiently basic or acidic to form stable nontoxic acid or base salts, administration of the compounds as salts may be appropriate. Examples of pharmaceutically acceptable salts are organic acid addition salts formed with acids which form a physiological acceptable anion, 5 for example, tosylate, methanesulfonate, acetate, citrate, malonate, tartarate, succinate, benzoate, ascorbate, α -ketoglutarate, and α -glycerophosphate. Suitable inorganic salts may also be formed, including hydrochloride, sulfate, nitrate, bicarbonate, and carbonate salts.

Pharmaceutically acceptable salts may be obtained using standard 10 procedures well known in the art, for example by reacting a sufficiently basic compound such as an amine with a suitable acid affording a physiologically acceptable anion. Alkali metal (for example, sodium, potassium or lithium) or alkaline earth metal (for example calcium) salts of carboxylic acids can also be made.

15 The compounds of the present invention can be formulated as pharmaceutical compositions and administered to a mammalian host, such as a human patient in a variety of forms adapted to the chosen route of administration, i.e., orally or parenterally (e.g., by intravenous, intramuscular, intraperitoneal). Preferably, the compounds are administered parenterally.

20 The active compound may also be administered intravenously or intraperitoneally by infusion or injection. Solutions of the active compound or its salts can be prepared in water, optionally mixed with a nontoxic surfactant. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, triacetin, and mixtures thereof and in oils. Under ordinary conditions of storage 25 and use, these preparations contain a preservative to prevent the growth of microorganisms.

The pharmaceutical dosage forms suitable for injection or infusion can include sterile aqueous solutions or dispersions or sterile powders comprising the active ingredient which are adapted for the extemporaneous 30 preparation of sterile injectable or infusible solutions or dispersions, optionally encapsulated in liposomes. In all cases, the ultimate dosage form should be sterile, fluid and stable under the conditions of manufacture and storage. The liquid carrier or vehicle can be a solvent or liquid dispersion medium

comprising, for example, water, ethanol, a polyol (for example, glycerol, propylene glycol, liquid polyethylene glycols, and the like), vegetable oils, nontoxic glyceryl esters, and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the formation of liposomes, by the maintenance
5 of the required particle size in the case of dispersions or by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, buffers or sodium chloride.

10 Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions are prepared by incorporating the active compound in the required amount in the appropriate solvent with various
15 of the other ingredients enumerated above, as required, followed by filter sterilization. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze drying techniques, which yield a powder of the active ingredient plus any additional desired ingredient present in the previously sterile-filtered
20 solutions.

For illustration, suitable doses of a compound of the invention for use in therapy, in conjunction with neutron capture, include doses in the range of from about 0.1 μ g to about 100 μ g, e.g., from about 0.5 μ g to about 50 μ g, or from about 0.5 μ g to 15 μ g per treatment. Suitable doses for use in imaging or
25 for use in imaging and therapy include doses in the range of from about 0.1 mg to about 50 g, e.g., from about .5 mg to about 10 g, or from about 0.5 g to 2 g per treatment.

The desired dose may conveniently be presented in a single dose or as divided doses administered at appropriate intervals, for example, as two,
30 three, four or more sub-doses per day. The sub-dose itself may be further divided, e.g., into a number of discrete loosely spaced administrations.

The compounds are preferably dissolved or dispersed in a nontoxic liquid vehicle, such as physiological saline or a similar aqueous

vehicle, to the desired concentration. A preselected therapeutic unit dose is then administered to the test animal or human patient, by oral administration or ingestion or by parenteral administration, as by intravenous or intraperitoneal infusion or injection, to attain the desired *in vivo* concentration. Doses useful for
5 treating tumors can be derived, from those found to be effective to treat tumors in humans *in vitro* or in animal models, such as those described hereinbelow, or from dosages of other labeled vitamin B₁₂ molecules, previously employed in animal therapy.

Figure 2 and Figure 3 each illustrate the four step synthesis used
10 to prepare three cyanocobalamin-nido-carborane conjugates. o-Carborane carboxylic acid (2) was prepared by reacting o-carborane with n-butyllithium and carbondioxide in ether for approximately one hour at -78°C. Treatment with thionyl chloride gave o-carborane carboxylic acid chloride (3), which was allowed to react with 1,4-butanediamine in pyridine to give the amide linked
15 nido-carboranoyl(4-aminobutyl)amide (4). Compound (4) was linked to the b-monocarboxylic acid of cyanocobalamin, the d-monocarboxylic acid of cyanocobalamin, or both the b- and d-dicarboxylic acid of cyanocobalamin. In each reaction, hydroxybenzotriazole and 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide was added to facilitate the formation of the amide bond.

20 The invention may be further illustrated by the following examples.

Examples

Ultraviolet-visible spectra were recorded with a diode array spectrophotometer. ¹¹B NMR spectra at 192.656 MHZ with ¹H decoupling were
25 recorded on Varian 1NOVA 600 MHZ spectrometer with an 8 mm broad band probe. Approximately 10 mg of the cobalamin-conjugate were dissolved in 1 mL pyridine d₆ in a 5 mm NMR tube. Five hundred scans were collected with an acquisition time of 0.133 seconds and a relaxation delay of 1 second. Chemical shifts are given relative to BF₃·(CH₃CH₂)₂O.

30 Mass spectral data was obtained on a Sciex API 365 LC-MS/MS system (Toronto, Canada). Separations were done on a Shimadzu HPLC system consisting of two LC-10AD pumps and a SCL-10Avp controller (Shimadzu Scientific Instruments, Columbia, MD). Analytes were monitored by UV at 214

NM with an ABI 785 detector. HPLC separations were achieved using a BDS-Hypersil C8 column (150 × 4.6 mm; 120 Å, Keystone Scientific, Inc., Bellefonte, PA). The mobile phase consisted of H₂O:methanol (98:2 v:v) in pump A and H₂O: methanol (2:92 v:v) in pump B. A linear gradient was used from 5% B to 30% B over 10 minutes and was held at 30% B for 20 minutes before returning to initial conditions (mono-carborane synthesis). The di-carborane required a longer gradient using the same mobile phases; 5% B to 65% B over 25 minutes and held at 65% B for 15 minutes before returning to initial conditions. The separations were monitored by UV absorption at 214 NM. The flow was 1.0 ml/minute and was split post-column allowing ~ 10 ul to flow into the mass spectrometer.

Mass spectral data was collected using electrospray ionization in positive mode over a mass range of 300 to 2300 AMU at a dwell time of 0.3 ms/0.1 AMU. Synthetic samples were prepared at 5 or 10 mg/ml in pump A mobile phase and an aliquot injected onto the HPLC (1-5 ul). Retention times of the mono- and di-carborane products were determined to be 13.3 minutes and 16.2 minutes respectively. Purification of the b and d mono-carborane products was achieved by collecting fractions at the elution times for several injections. The collected fractions were combined and dried to a powder. A portion of the purified product was dissolved in methanol:H₂O (1:1) and reanalyzed by HPLC-MS to ascertain the purity.

α-Carborane, butyllithium (1.6 M solution in hexanes) and putrescine were purchased from Aldrich Chemical Company (Milwaukee, WI). The water-soluble carbodiimide 1-ethyl-3-(3'-dimethylaminopropyl) carbodiimide and 1-hydroxybenzotriazole were from Sigma (St. Louis, MO). Thin layer chromatography (TLC) silica gel plates were obtained from Eastman Kodak Company. The cyanocobalamin-b, d, and e monocarboxylic acids and the b, d-dicarboxylic acid were prepared as described before (see U.S. Patent Number 5,739,313, and D.L. Anton, et al., *J. Am. Chem. Soc.*, 1980, 102, 2212-2219), and were provided by Professor Kahl of the Department of Medicinal Chemistry at the University of California, San Francisco.

Example 1 Cyanocobalamin-nido-carborane conjugates (5, Figure 2).

Separate reaction mixtures containing 1 g (~0.66 mmol) of the b-monocarboxylic acid, d-monocarboxylic acid, or the b,d-dicarboxylic acid of cyanocobalamin, hydroxybenzotriazole (810 mg, 6.0 mmol), 1-ethyl-3(3-
 5 dimethylaminopropyl) carbodiimide (11.4 g, 6.0 mmol) and 4 (600 mg, 2.0 mmol) in 100 mL of a water-acetone mixture (2:1) were adjusted to pH 6.9 with 1N NaOH. The reactions were stirred at room temperature and their progress monitored by TLC. After 3 hours, the mixtures were concentrated to remove acetone and the resulting aqueous suspensions were extracted into 92% aqueous
 10 phenol. The phenol phases were washed with water to remove the water-soluble reactants. One volume of acetone and three volumes of ether were added to each of the phenol phases and the desired cyanocobalamin-nido-carborane conjugates were back extracted into water. The aqueous layers were extracted three times with ether to remove residual phenol and unreacted (4). Finally the aqueous
 15 solutions were evaporated to dryness. The residuals were triturated with acetone and the desired conjugates isolated as orange-red powders (yields 90-95% based on the cyanocobalamin-carboxylic acids).

The isotope ratios of the final products were also compared to theoretical as follows: Mono product theoretical $M+H^+$
 20 (C₇₀H₁₀₉CoB₉O₁₅PN₁₅) - 1584.8 (12%), 1585.8 (37%), 1586.8 (74%), 1587.8 (100%), 1588.8 (84%), 1589.8 (44%), 1590.8 (16%), 1591.8 (4%). Synthesis I - 1584.8 (14%), 1585.8 (37%), 1586.8 (74%), 1587.8 (100%), 1588.8 (88%), 1589.8 (48%), 1590.8 (19%), 1591.8 (6%). Synthesis II - 1584.8 (15%), 1585.8 (40%), 1586.8 (78%), 1587.8 (100%), 1588.8 (86%), 1589.8 (44%), 1590.8
 25 (17%), 1591.8 (7%).

Di product theoretical $M+H^+$ - (C₇₇H₁₂₉CoB₁₈O₁₆PN₁₆) - 1815.1 (9%), 1816.1 (23%), 1817.1 (47%), 1818.1 (77%), 1819.1 (99%), 1820.1 (100%), 1821.1 (77%), 1822.1 (44%), 1823.1 (19%), 1824.1 (6%). Observed synthesis - 1815.1 (15%), 1816.1 (27%), 1817.1 (51%), 1818.1 (79%), 1819.1
 30 (100%), 1820.1 (100%), 1821.1 (78%), 1822.1 (49%), 1823.1 (25%), 1824.1 (12%).

Results included the identification of the *b* and *d* carborane cyanocobalamin (CCC) analogs by LC-MS. Products identified were

subsequently separated and purified by HPLC. The purified products were reanalyzed by LC-MS with no starting material detected. MS/MS data was also obtained on the *b* and *d* CCC analogs to provide further structural characterization. These purified products were then tested in the biological assays. The *e*-CCC analog and the *di*-CCC analog were also separated and identified by LC-MS, however, the preparations contained more reaction side products resulting in difficult purification. The isotope ratios of the *b*- and *d*-CCC analogs, as well as the *di*-CCC analog, were compared to theoretical isotope ratios to provide further information for identification.

Mass spectra confirmed the presence of each cyanocobalamin-*nido*-carborane conjugate. Mass spectroscopy analysis as well as ^{11}B NMR showed that during the conversion of 3 to 4 (Figures 2 and 3) the *o*-carborane nucleus lost a boron atom to yield the *nido*-carborane derivative 4 (Figure 3). UV-visible spectroscopy of the final products showed maxima typical of cyanocobalamin, indicating that the corrin nucleus was intact and that the 5,6-dimethyl benzimidazole nucleotide was still attached to the cobalt atom.

The starting material 4 was prepared as follows (see Figure 2):

a. *o*-carborane carboxylic acid (2). A solution of *o*-carborane (1) (5.0 g, 34.7 mmol) in 500 ml dry ether in a 1 L round bottom flask was cooled to -78°C in a dry ice-acetone bath. The solution was flushed with argon and the flask sealed with a serum stopper. *n*-Butyllithium (24 mL, 1.6 M in hexanes) was slowly injected via a syringe over a period of about 20 minutes and the reaction was stirred for an additional 30 minutes. Crushed dry ice (10-15 g) was then added and the mixture stirred for 1 hour. The dry ice-acetone bath was removed and the reaction allowed to come to room temperature. The ether and hexanes were removed on a rotary evaporator, water (150 ml) was added and unreacted *o*-carborane was removed by extraction with hexanes (2×100 ml). The aqueous phase was acidified with concentrated HCl and the desired product extracted with hexanes (4×100 ml). The combined extracts were dried over Na_2SO_4 and evaporated to dryness to give 5.8 g (88.7%) of *o*-carborane carboxylic acid (2), which was used without further purification.

b. o-carborane carboxylic acidchloride (3). o-Carborane carboxylic acid (2.0 g, 10.6 mmol), dried over P_2O_5 , was dissolved in 30 ml thionylchloride and heated under reflux for 3 hours. The solution was cooled to room temperature, evaporated to dryness and dried over P_2O_5 (2.05 g, 9.9 mmol, 5 93%) to provide (3), which was used without further purification.

c. Nido-Carboranoyl(4-aminobutyl) amide (4). The acid chloride (3) was dissolved in 15 ml dry pyridine and 1,4-diaminobutane (1.12 g, 12.7 mmol) was added. The reaction mixture was heated under reflux for 3 hours, cooled to room temperature and concentrated. Water (10 ml) was added 10 and the suspension acidified with 3 M HCl. The desired product was extracted with ethylacetate (4×50 ml), the combined organic layers were washed once with water, dried over Na_2SO_4 and evaporated to dryness to yield 2.35 g (8.0 mmol, 75%) of (4). Thus far the product has resisted crystallization from a variety of solvent mixtures, however TLC on silica gel plates (2-propanol- 15 $NH_4OH \cdot H_2O$; 7:1:2) showed only one ninhydrin positive compound distinct from the diamine.

Example 2. 6-Carboranoylamidopropyl cobalamin.

6-(3-aminoprop-1-yl)cobalamin (300 mg), hydroxybenzotriazole (270 mg), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (382 mg) and o- 20 carborane monocarboxylic acid were dissolved in water (50 ml) and acetone (20 ml) and allowed to react at room temperature for 3 hours. The reaction mixture was concentrated to remove acetone and desalted via phenol extraction. The concentrated aqueous solution was crystallized from aqueous acetone to give the title compound.

25 The intermediate 6-(3-aminoprop-1-yl)cobalamin was prepared as follows:

a. 6-(3-aminoprop-1-yl)cobalamin. Cyano cobalamin (500 mg) was dissolved in 100 ml deoxygenated water containing 10 mg $CoCl_2$, and reduced with sodium borohydride to give the corresponding Co (I) compound. 30 After 30 minutes, 3-chloropropyl amine hydrochloride (130 mg) dissolved in 5 ml deoxygenated ethanol was added. After one hour, at room temperature, the mixture was desalted via phenol extraction. The aminopropylcarbolamine was back extracted into water after the addition of 1 volume of acetone and 3

volumes of ether. The aqueous solution was concentrated and the desired 6-(3-aminoprop-1-yl)cobalamin crystallized from aqueous acetone (yield 510 mg).

Example 3 DTPA-aminopropylcobalamin (DAPC)

5 Using a coupling procedure similar to that described in Example 2, 6-(3-aminoprop-1-yl)cobalamin and DTPA were coupled in the presence of hydroxybenzotriazole to give the title compound.

Example 4 *In vitro* biological activity of the carborane cyanocobalamin analogs.

10 To assess the *in vitro* binding of the carborane cyanocobalamin (Example 1, CCC) and DTPA-aminopropylcobalamin (Example 3, DAPC) analogs to the transcobalamin proteins, the unsaturated Vitamin B12 binding capacity (UBBC) assay (see D.A. Collins and H.P.C. Hogenkamp, *J. Nuclear Medicine*, 1997, 38, 717-723) was performed. Serum was obtained from 5
15 patients being evaluated for pernicious anemia at the Mayo Clinic. The patients' serum first underwent a routine clinical UBBC as previously described. To determine if the CCC and DAPC analogs would inhibit Co-57 cyanocobalamin from binding to the transcobalamin proteins, the excess serum from the 5 patients underwent modified UBBC.

20 Specifically, under dim light, 0.4 ml serum was treated with 4 μ l (concentration 10 μ g CCC per ml normal saline) of the 2 CCC analogs, carborane-d-cyanocobalamin and carborane-b-cyanocobalamin, and DAPC. The analogs were incubated for 20 minutes at room temperature with the patient's serum. Both the clinical run and the analog treated samples were assayed for
25 UBBC as usual.

The analogs competitively blocked Co-57-cyanocobalamin from binding to the transcobalamin proteins. Therefore, the cpm of the modified UBBC assay was significantly lower than that of the clinical runs. The percent PB of the analogs to transcobalamin protein was calculated as follows (PB = 100
30 - CCC_{UBBC} cpm/clinical UBBC cpm \times 100). The average percent binding (PB) of the 5 solutions (n = 10 for each solution, i.e., two modified UBBC assays per patient) for the carborane-d cyanocobalamin and carborane-b cyanocobalamin was 35.75% and 92.93% respectively, and 98.21% for DAPC.

All publications, patents, and patent documents are incorporated by reference herein, as though individually incorporated by reference. The invention has been described with reference to various specific and preferred embodiments and techniques. However, it should be understood that many
5 variations and modifications may be made while remaining within the spirit and scope of the invention.

What is claimed is:

1. A compound wherein a residue of a compound of formula I (Figure 1) is linked to a residue of a molecule comprising B-10, wherein X is
5 CN, OH, CH₃, adenosyl, or a molecule comprising B-10; or a pharmaceutically acceptable salt thereof.
2. The compound of claim 1 wherein the residue of a molecule comprising B-10 is directly linked to the 6-position of the compound of formula
10 I or is directly linked to a residue of the b, d or e-carboxamide group of the compound of formula I.
3. The compound of claim 1 wherein the residue of a molecule comprising B-10 is linked by a linker to the 6-position of the compound of
15 formula I or is linked by a linker to the residue of a-, b-, d- or e-carboxamide group of the compound of formula I.
4. The compound of claim 1 wherein the residue of a molecule comprising B-10 is linked to a residue of the b-carboxamide group of the
20 compound of formula I.
5. The compound of claim 1 wherein the residue of a molecule comprising B-10 is linked to a residue of d-carboxamide group of the compound
of formula I.
25
6. The compound of claim 1 wherein the residue of a molecule comprising B-10 is linked to a residue of the e-carboxamide group of the compound of formula I.
- 30 7. The compound of claim 1 wherein a residue of a molecule comprising B-10 is linked to a residue of the b-carboxamide group and a second residue of a molecule comprising B-10 is linked to a residue of the d-carboxamide group of the compound of formula I.

8. The compound of claim 1 wherein the residue of a molecule comprising B-10 is linked to the 6-position of the compound of formula I.
9. The compound of claim 1 wherein the molecule comprising B-10 contains 1 to about 20 boron atoms, inclusive.
10. The compound of claim 1 wherein the molecule comprising B-10 is an amino acid, a carbohydrate, a nucleoside or a carborane.
- 10 11. The compound of claim 1 wherein the molecule comprising B-10 is o-carborane, m-carborane or p-carborane.
12. The compound of claim 1 wherein the molecule comprising B-10 is o-carborane.
- 15 13. The compound of claim 3 wherein at least one linker is of the formula W-A-Q wherein A is (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, (C₃-C₈)cycloalkyl, or (C₆-C₁₀)aryl, wherein W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)₂-, -N(R)-, -C(=O)-, or a direct bond; wherein each R is independently H or (C₁-C₆)alkyl.
- 20 14. The compound of claim 13 wherein W is NH₂ or COOH and Q is NH₂ or COOH.
- 25 15. The compound of claim 13 wherein A is (C₁-C₆)alkyl.
16. The compound of claim 3 wherein at least one linker is about 5 angstroms to about 50 angstroms, inclusive.
- 30 17. The compound of claim 3 wherein at least one linker comprises a therapeutic radionuclide or a diagnostic radionuclide.

18. The compound of claim 17 wherein the therapeutic radionuclide is a metallic radionuclide.
19. The compound of claim 17 wherein the diagnostic radionuclide is
5 a metallic radionuclide.
20. The compound of claim 17 wherein the diagnostic radionuclide is a non-metallic radionuclide.
- 10 21. The compound of claim 3 wherein at least one linker is a divalent radical formed from a peptide.
22. The compound of claim 3 wherein at least one linker is a divalent radical formed from an amino acid.
- 15 23. The compound of claim 3 wherein at least one linker is poly-L-glutamic acid, poly-L-aspartic acid, poly-L-histidine, poly-L-ornithine, poly-L-serine, poly-L-threonine, poly-L-tyrosine, poly-L-lysine-L-phenylalanine, poly-L-lysine or poly-L-lysine-L-tyrosine.
- 20 24. The compound of claim 1 wherein the residue of the compound of formula I is also linked to a linker comprising a detectable radionuclide or a therapeutic radionuclide.
- 25 25. A compound wherein a residue of a compound of formula I (Figure 1) is linked to a group of the formula Q-L-W-Det; wherein X is CN, OH, CH₃, adenosyl, a molecule comprising B-10, or Q-L-W-Det; wherein Det is a chelating group comprising Gd-157; L is a linker or absent; and W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-,
30 -S(O)-, -S(O)₂-, -C(=O)-, -N(R)-, or a direct bond; wherein each R is independently H or (C₁-C₆)alkyl; or a pharmaceutically acceptable salt thereof.

26. The compound of claim 25 wherein the group of the formula Q-L-W-Det is linked to a residue of the b-carboxamide group, a residue of the d-carboxamide group, a residue of the e-carboxamide, or the 6-position of the compound of formula I.
- 5 27. The compound of claim 25 wherein the group of the formula Q-L-W-Det is linked to a residue of the b-carboxamide group and a second group of the formula Q-L-W-Det is linked to a residue of the d-carboxamide group of the compound of formula I.
- 10 28. The compound of claim 25 wherein the group of the formula Q-L-W-Det is between about 20 and about 500 angstroms, inclusive, in length.
29. The compound of claim 25 wherein at least one chelating group is
15 EDTA, DTPA, DOTA, DOTMP, TETA, MAG3 or DCTA.
30. The compound of claim 25 wherein at least one chelating group is DTPA comprising Gd-157.
- 20 31. A compound wherein a residue of a compound of formula I (Figure 1) is linked to a residue of a molecule comprising B-10; wherein a residue of the compound of formula I is also linked to a group of the formula Q-L-W-Det, wherein X is CN, OH, CH₃, adenosyl, a group of the formula Q-L-W-Det, or a molecule comprising B-10; wherein:
- 25 Det is a chelating group comprising a therapeutic radionuclide or a diagnostic radionuclide;
- L is a linker or absent; and
- Q and W are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)₂-, -C(=O)-, -N(R)-, or a direct
30 bond; wherein each R is independently H or (C₁-C₆)alkyl; or a pharmaceutically acceptable salt thereof.

32. The compound of claim 31 wherein at least one of the radionuclides is Tc^{99m}, In¹¹¹, In¹¹⁰, Gd¹⁵⁷ or Y⁸⁶.
33. The compound of claim 31 wherein a molecule comprising B-10 is linked to the residue of a b-carboxamide group, d-carboxamide group, e-carboxamide group, or the 6-position of the compound of formula I.
34. The compound of claim 31 wherein the chelating group is DTA, DTPA, DOTA, TETA, DOTMP, MAG3 or DCTA.
35. The compound of claim 31 wherein the chelating group is DTPA.
36. The compound of claim 31 wherein the residue of a molecule comprising B-10 contains 1 to about 20 boron atoms.
37. The compound of claim 31 wherein the molecule comprising B-10 is a carbohydrate, an amino acid, a nucleoside or a carborane.
38. The compound of claim 31 wherein the molecule comprising B-10 is o-nido-carborane, m-nido-carborane or p-nido-carborane.
39. The compound of claim 31 wherein the molecule comprising B-10 is o-carborane.
40. The compound of claim 31 wherein the residue of a molecule comprising B-10 is directly linked to the 6-position or to the residue of the b-, d- or e-carboxamide group of the compound of formula I.
41. The compound of claim 31 wherein the residue of the compound of formula I is linked to a residue of a molecule comprising B-10 through a linker.

42. The compound of claim 41 wherein the linker comprises a non-metallic radionuclide.
43. The compound of claim 41 wherein the linker is about 5
5 angstroms to about 50 angstroms, inclusive.
44. The compound of claim 1 further comprising a detectable radionuclide.
- 10 45. The compound of claim 2 wherein the detectable radionuclide is a non-metallic radionuclide.
46. The compound of claim 3 wherein the non-metallic radionuclide is Carbon-11, Fluorine-18, Bromine-76, Iodine-123, or Iodine-124.
- 15 47. The compound of claim 2 wherein the detectable radionuclide is directly linked to the compound of formula I.
48. The compound of claim 2 wherein the detectable radionuclide is
20 linked by a linker to the compound of formula I.
49. The compound of claim 6 wherein the linker is of the formula W-A wherein A is (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, (C₃-C₈)cycloalkyl, or (C₆-C₁₀)aryl, wherein W is -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-,
25 -O-, -S-, -S(O)-, -S(O)₂-, -N(R)-, -C(=O)-, or a direct bond; wherein each R is independently H or (C₁-C₆)alkyl; and wherein A is substituted with one or more non-metallic radionuclides.
50. The compound of claim 48 wherein the linker is about 5
30 angstroms to about 50 angstroms, inclusive.
51. The compound of claim 48 wherein the linker is a divalent peptide or amino acid.

52. The compound of claim 48 wherein the linker is poly-L-glutamic acid, poly-L-aspartic acid, poly-L-histidine, poly-L-ornithine, poly-L-serine, poly-L-threonine, poly-L-tyrosine, poly-L-lysine-L-phenylalanine, or poly-L-lysine or poly-L-lysine-L-tyrosine.
53. The compound of claim 48 wherein the linker is linked to the 6-position of the compound of formula I or is linked to the residue of a-, b-, d- or e-carboxamide group of the compound of formula I.
54. A compound wherein a residue of a compound of formula I (Figure 1) is linked 1) to a detectable radionuclide; and is linked 2) to a group comprising Gd-157; or a pharmaceutically acceptable salt thereof.
55. The compound of claim 54 wherein the group comprising Gd-157 has the formula Q-L-W-Det; wherein X is CN, OH, CH₃, adenosyl, a molecule comprising B-10, or Q-L-W-Det; Det is a chelating group comprising Gd-157; L is a linker or absent; and W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)₂-, -C(=O)-, -N(R)-, or a direct bond; wherein each R is independently H or (C₁-C₆)alkyl.
56. The compound of claim 54 wherein the detectable radionuclide is a non-metallic radionuclide.
57. The compound of claim 56 wherein the non-metallic radionuclide is Carbon-11, Fluorine-18, Bromine-76, Iodine-123, or Iodine-124.
58. The compound of claim 54 wherein the detectable radionuclide is directly linked to the compound of formula I.
59. The compound of claim 54 wherein the detectable radionuclide is linked by a linker to the compound of formula I.

60. The compound of claim 59 wherein the linker is of the formula W-A wherein A is (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, (C₃-C₈)cycloalkyl, or (C₆-C₁₀)aryl, wherein W is -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)₂-, -N(R)-, -C(=O)-, or a direct bond; wherein each R is independently H or (C₁-C₆)alkyl; and wherein A is substituted with one or more non-metallic radionuclides.
61. The compound of claim 59 wherein the linker is about 5 angstroms to about 50 angstroms, inclusive.
62. The compound of claim 59 wherein the linker is a divalent peptide or amino acid.
63. The compound of claim 59 wherein the linker is poly-L-glutamic acid, poly-L-aspartic acid, poly-L-histidine, poly-L-ornithine, poly-L-serine, poly-L-threonine, poly-L-tyrosine, poly-L-lysine-L-phenylalanine, or poly-L-lysine or poly-L-lysine-L-tyrosine.
64. The compound of claim 59 wherein the linker is linked to the 6-position of the compound of formula I or is linked to the residue of a-, b-, d- or e-carboxamide group of the compound of formula I.
65. A compound wherein a residue of a compound of formula I (Figure 1) is linked 1) to a molecule comprising B-10 or to a chelating group comprising Gd-157, and 2) to at least one residue of the formula Q-L-W-Det; wherein each Det is independently a chelating group comprising a metallic radionuclide; each L is independently a linker or absent; and each W and Q is independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)₂-, -C(=O)-, -N(R)-, or a direct bond; wherein each R is independently H or (C₁-C₆)alkyl; or a pharmaceutically acceptable salt thereof.
66. The compound of claim 1 or 44 wherein the residue of a compound of formula I is also linked to a group comprising Gd-157.

67. The compound of claim 66 wherein the group comprising Gd-157 has the formula Q-L-W-Det; wherein X is CN, OH, CH₃, adenosyl, a molecule comprising B-10, or Q-L-W-Det; Det is a chelating group comprising Gd-157; L is a linker or absent; and W and Q are each independently -N(R)C(=O)-, -C(=O)N(R)-, -OC(=O)-, -C(=O)O-, -O-, -S-, -S(O)-, -S(O)₂-, -C(=O)-, -N(R)-, or a direct bond; wherein each R is independently H or (C₁-C₆)alkyl.
68. A pharmaceutical composition comprising a compound of any one of claims 1-67 and a pharmaceutically acceptable carrier.
69. A method of treating a tumor in a mammal in need of such treatment comprising administering to the mammal an effective amount of a compound of any one of claims 1-67 in combination with a pharmaceutically acceptable vehicle; and administering neutron capture therapy.
70. A method for imaging a tumor in a mammal comprising administering to the mammal a detectable amount of a compound of any one of claims 1-67; and detecting the presence of the compound.
71. The method of claim 70 further comprising treating the tumor with neutron capture therapy.
72. A compound of any one of claims 1-67 for use in medical therapy or diagnosis.
73. The use of a compound of any one of claims 1-67 for the manufacture of a medicament for imaging a tumor in a mammal.
74. The use of a compound of any one of claims 1-67 for the manufacture of a medicament for treating a tumor in a mammal in need of such treatment.





FIG. 1

2/3



(o-CARB) = o-Carborane

$[\text{o-nido-CARB}]^-$ = o-nido-Carborane

FIG. 2




FIG. 3

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 October 2000 (26.10.2000)

PCT

(10) International Publication Number
WO 00/62808 A3

(51) International Patent Classification⁷: A61K 41/00

US 60/159,873 (CIP)
Filed on 15 October 1999 (15.10.1999)

(21) International Application Number: PCT/US00/10100

(22) International Filing Date: 15 April 2000 (15.04.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/129,733 16 April 1999 (16.04.1999) US
60/159,873 15 October 1999 (15.10.1999) US


(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:
US 60/129,733 (CIP)
Filed on 16 April 1999 (16.04.1999)

(71) Applicants (for all designated States except US): MAYO OF FOUNDATION FOR MEDICAL EDUCATION AND RESEARCH [US/US]; 200 First Street S.W., Rochester, MN 55905 (US). REGENTS OF THE UNIVERSITY OF MINNESOTA [US/US]; 600 University Gateway, 200 Oak Street SE, Minneapolis, MN 55455-2070 (US).

(71) Applicants and
(72) Inventors: COLLINS, Douglas, A. [US/US]; 1150 Meadowlark Court S.W., Rochester, MN 55902 (US). HOGENKAMP, Henricus, P., C. [US/US]; 2211 Marion Road, Roseville, MN 55113 (US).

[Continued on next page]

(54) Title: COBALAMIN CONJUGATES USEFUL AS ANTITUMOR AGENTS



[Continued on next page]



(74) Agent: VIKSNINS, Ann, S.; Schwegman, Lundberg, Woessner & Kluth, P.O. Box 2938, Minneapolis, MN 55402 (US).

(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

Published:

— With international search report.

(88) Date of publication of the international search report:
22 February 2001

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(57) Abstract: The invention provides cobalamin compounds linked to a neutron capture therapy target (e.g. Boron-10 or Gadolinium-157), and optionally linked to a detectable moiety, as well as pharmaceutical compositions comprising the compounds, and methods for using the compounds in medical diagnosis and therapy.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 00/10100

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K41/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EMBASE, BIOSIS, EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	HOGENKAMP, H. P. C. ET AL: "Synthesis and characterization of nido-carborane-cobalamin conjugates" NUCL. MED. BIOL. (2000), 27(1), 89-92 , XP002148452 cited in the application page 89, column 1, paragraphs 1,2; figures 1,2	1-74
E	WO 00 45857 A (SCHERING AG) 10 August 2000 (2000-08-10) claims	1-74
A	WO 92 09160 A (TAU SYSTEMS CORP) 29 May 1992 (1992-05-29) claims	
	-/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

26 September 2000

Date of mailing of the international search report

13/10/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Berte, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/10100

C.,(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 739 313 A (COLLINS DOUGLAS A ET AL) 14 April 1998 (1998-04-14) cited in the application column 3, line 44 - line 53; claims column 4, line 36 - line 40 -----	1
A	US 5 877 165 A (MIURA MICHIKO ET AL) 2 March 1999 (1999-03-02) claims -----	1-74

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claim(s) 70 are directed to a diagnostic method practised on the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Although claims 69,71 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

INTERNATIONAL SEARCH REPORT

Information on patent family members

In. ational Application No

PCT/US 00/10100

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0045857 A	10-08-2000	NONE	
WO 9209160 A	29-05-1992	US 5103476 A US 5222134 A AT 171024 T CA 2095723 A DE 69130175 D DE 69130175 T EP 0556305 A JP 7089345 B JP 6501120 T	07-04-1992 22-06-1993 15-09-1998 08-05-1992 15-10-1998 10-02-2000 25-08-1993 27-09-1995 27-01-1994
US 5739313 A	14-04-1998	AU 702026 B AU 1119597 A AU 719545 B AU 2499999 A CA 2237263 A EP 0883627 A JP 2000500753 T US 6004533 A WO 9718231 A US 6096290 A	11-02-1999 05-06-1997 11-05-2000 17-06-1999 22-05-1997 16-12-1998 25-01-2000 21-12-1999 22-05-1997 01-08-2000
US 5877165 A	02-03-1999	NONE	

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 October 2000 (26.10.2000)

PCT

(10) International Publication Number
WO 00/62808 A3

(51) International Patent Classification⁷: A61K 41/00

US

60/159,873 (CIP)

(21) International Application Number: PCT/US00/10100

Filed on

15 October 1999 (15.10.1999)

(22) International Filing Date: 15 April 2000 (15.04.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/129,733 16 April 1999 (16.04.1999) US

60/159,873 15 October 1999 (15.10.1999) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:

US 60/129,733 (CIP)

Filed on 16 April 1999 (16.04.1999)


(71) Applicants (for all designated States except US): MAYO FOUNDATION FOR MEDICAL EDUCATION AND RESEARCH [US/US]; 200 First Street S.W., Rochester, MN 55905 (US). REGENTS OF THE UNIVERSITY OF MINNESOTA [US/US]; 600 University Gateway, 200 Oak Street SE, Minneapolis, MN 55455-2070 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): COLLINS, Douglas, A. [US/US]; 1150 Meadowlark Court S.W., Rochester, MN 55902 (US). HOGINKAMP, Henricus, P., C. [US/US]; 2211 Marion Road, Roseville, MN 55113 (US).

[Continued on next page]

(54) Title: COBALAMIN CONJUGATES USEFUL AS ANTITUMOR AGENTS



(57) Abstract: The invention provides cobalamin compounds linked to a neutron capture therapy target (e.g. Boron-10 or Gadolinium-157), and optionally linked to a detectable moiety, as well as pharmaceutical compositions comprising the compounds, and methods for using the compounds in medical diagnosis and therapy.

WO 00/62808 A3



(74) **Agent:** KNOWLES, Sherry, M.: KING & SPALDING,
191 Peachtree Street, N.E., Atlanta, GA 30303-1763 (US).

Published:

— with international search report

(81) **Designated States (national):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(88) **Date of publication of the international search report:**
22 February 2001

(48) **Date of publication of this corrected version:**
6 December 2001

(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(15) Information about Correction:

see PCT Gazette No. 49/2001 of 6 December 2001, Section II

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.